

Empreinte sur l'Eau et le Bassin du Nil

Module 4: Analyse de l'empreinte sur
l'eau & méthodologie



PEGASYS
Strategy and Development





Aperçu de ce module

1^{ère} Section : Analyse et méthodologie de l'empreinte sur l'eau

3^{ème} Section : Sources des données et outils utiles

4^{ème} Section : Utilisant CROPWAT



Analyse de l'empreinte sur l'eau & méthodologie

Comprendre les indicateurs





Structure de cette séance

1. Empreinte sur l'eau d'un produit
2. Exemple détaillé: empreinte sur l'eau d'un produit cultivé
3. Exemple détaillé: empreinte sur l'eau d'un produit de bétail
4. Empreinte sur l'eau d'un pays



Empreinte sur l'eau

- Le volume d'eau fraîche nécessaire à la production du produit, reflétant le volume d'eau utilisée aux différentes étapes de la chaîne de production.
- Où et quand l'eau fut utilisée : une empreinte sur l'eau inclut une dimension temporelle et spatiale.
Tel que :
 - Climat (évapotranspiration)
 - Précipitations
 - Productivité



Eléments de l'empreinte sur l'eau

Empreinte sur l'eau verte – volume d'humidité du sol (eau de pluie) évaporée au cours de la croissance d'une culture


Empreinte sur l'eau bleue – volume de surface irriguée ou d'eau de surface évaporées ou incorporées dans un produit

Empreinte sur l'eau grise – volume d'eau nécessaire pour diluer la contamination de l'eau par le déversement de déchets et le retour d'écoulement





Exemple d'empreinte sur l'eau : Bière





Exemple détaillé: Empreinte sur l'eau d'un produit cultivé

E sur E =

$$\frac{\text{Eau nécessaire à une culture (m}^3/\text{ha})}{\text{Rendement d'une culture (tonne/ha)}}$$



Exemple détaillé: Empreinte sur l'eau d'un produit cultillé

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture verte} + \text{Eau nécessaire à une culture bleue}}{\text{Rendement d'une culture (tonne/ha)}}$$




Exemple détaillé: Empreinte sur l'eau d'un produit cultivé

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture}_{\text{verte}} + \text{Eau nécessaire à une culture}_{\text{bleue}}}{\text{Rendement d'une culture (tonne/ha)}}$$

Qu'est-ce que
l'eau verte
nécessaire à une
culture ?



Empreinte sur l'eau d'un produit cultivé – Eau verte






Exemple détaillé: Empreinte sur l'eau d'un produit cultivé

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture}_{\text{verte}} + \text{Eau nécessaire à une culture}_{\text{bleue}}}{\text{Rendement d'une culture (tonne/ha)}}$$

Qu'est-ce que
l'eau bleue
nécessaire à une
culture ?



Empreinte sur l'eau d'un produit cultivé – Eau bleue



$$E \text{ sur } E_{\text{bleue}} = \text{Evapo-} \\ \text{transpiration} + \text{Incorporation} \\ \text{d'eau bleue dans} \\ \text{une culture}$$



Méthodes d'évaluation CROPWAT

1. Option CWR (*crop water requirements; besoins en eau d'une culture*)

- Conditions optimales
- Sans considération pour la balance d'eau de sol
- Intervalles de temps de 10 jours entre précipitations réelles

2. Option programme d'irrigation – Pas de conditions optimales

- Considération pour la balance d'eau de sol
- Intervalles de temps quotidiennes pour la balance d'eau de sol




1. Option CWR

- Données climatiques

Données climatiques – base de données **CLIMWAT 2.0** (FAO, 2006)

Données agroclimatiques observées dans plus de 5000 stations à travers le monde (tentative de couvrir la période 1971-2000)






1. Option CWR

- Evapotranspiration des cultures

FIGURE 4. Reference (ET_0), crop evapotranspiration under standard (ET_C) and non-standard conditions ($ET_{C\ adj}$)



Allen et al. (1998)

Coefficient d'une culture

Caractéristiques: Hauteur de la culture, albédo (facteur de réflexion), résistance de voûte, évaporation du sol.

Evapotranspiration d'une culture de référence

Taux d'évapotranspiration d'une surface d'herbe de référence hypothétique qui est suffisamment irriguée


Evapotranspiration d'une culture dans des conditions normales

Evapotranspiration de cultures exemptes de maladies, convenablement fertilisées, cultivées dans de larges champs, avec des conditions d'eau de sol optimales et débouchant sur une production complète dans des conditions climatiques données



1. Option CWR


- Courbe de coefficient d'une culture






1. Option CWR

- Etapes de croissance et de développement de différentes cultures





Exemple détaillé : Utilisation d'eau verte





Exemple détaillé : Empreinte d'une culture sur l'eau verte

Evapotranspiration verte =


min(évapotranspiration d'une culture, précipitations réelles)

Eau nécessaire à une culture verte = $10 \times \sum$ Evapotranspiration verte

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture (m}^3/\text{ha})}{\text{Rendement d'une culture (tonne/ha)}}$$



Exemple détaillé : Utilisation d'eau bleue





Exemple détaillé : Empreinte d'une culture sur l'eau bleue

Evapotranspiration bleue =




min(besoins d'irrigation d'une culture, irrigation réelle)

Eau nécessaire à une culture bleue = $10 \times \sum$ Evapotranspiration bleue

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture (m}^3/\text{ha})}{\text{Rendement d'une culture (tonne/ha)}}$$




Données de CROPWAT





Exemple détaillé : Empreinte sur l'eau d'une culture

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture (m}^3/\text{ha})}{\text{Rendement d'une culture (tonne/ha)}}$$


Rendement d'une culture =
production d'une culture par
unité de terre cultivée




Qu'est-ce qui influence l'empreinte sur l'eau ?

$$E \text{ sur } E = \frac{\text{Eau nécessaire à une culture (m}^3/\text{ha})}{\text{Rendement d'une culture (tonne/ha)}}$$

Type de culture / produit
Evapotranspiration d'une culture : climat




Utilisation de fertilisants
Pratiques d'irrigation
Pas de réception d'eau optimale
Sol





Empreinte sur l'eau d'une culture – eau grise



- Volume d'eau fraîche polluée par la production d'une culture, telle que l'utilisation de fertilisants, de produits chimiques, et du niveau de salinité.
- Calculé grâce au volume d'eau nécessaire pour assimiler les polluants, sur base des normes de qualité de l'eau ambiante.




Exemple détaillé : Empreinte sur l'eau d'un produit de bétail





Exemple d'empreinte sur l'eau : un animal vivant





Empreinte sur l'eau d'un animal vivant

$$E \text{ sur } E_{\text{animal vivant}} = E \text{ sur } E_{\text{bue}} + E \text{ sur } E_{\text{entretien}} \\ + E \text{ sur } E_{\text{alimentation}}$$

- $E \text{ sur } E_{\text{bue}}$ = volume d'eau bu par un animal
- $E \text{ sur } E_{\text{entretien}}$ = volume d'eau nécessaire au maintien de l'animal
- $E \text{ sur } E_{\text{alimentation}}$ = eau contenue dans les cultures nécessaires à la production de l'alimentation pour le bétail



Empreinte sur l'eau d'un animal vivant

- Empreinte sur l'eau de son alimentation



- Empreinte sur l'eau de l'herbe (m³/tonne)
- Volume d'herbe mangée par l'animal (tonne)

$$E_{\text{sur E}} = E_{\text{Herbe}} * \text{Volume}_{\text{Herbe mangée}}$$




Empreinte sur l'eau d'un pays






Empreinte sur l'eau d'un pays






Cadre de comptabilisation de l'eau utilisée au niveau national





Importateurs et exportateurs nets d'eau



Balances nationales d'eau virtuelle liées au commerce international de produits. Période 1997-2001. Exportateurs nets marqués en vert et importateurs nets en rouge.



Sources des données et outils utiles





Analyse de l'empreinte sur l'eau

Données & information

Données requises :

- Données sur les précipitations et le taux d'évaporation
- Données sur la croissance des cultures et l'évapotranspiration
- Pratiques d'irrigation
- Rendement des cultures
- Volumes de production
- Volumes d'échanges commerciaux
- Partenaires commerciaux

Informations supplémentaires :

- Emplois associés à la production
- Contribution au PIB
- Volumes de relations commerciales informelles (si pertinent et disponible)



Analyse de l'empreinte sur l'eau

Données & information

Sources d'informations utiles et des outils:



Les précipitations et d'évaporation

<http://www.fao.org/nr/water/infore databases climwat.html>

► text only version ► print friendly

helping to build a world without hunger

NATURAL RESOURCES AND ENVIRONMENT DEPARTMENT



FAO
WATER

FrontPage

Topics

Information Resources

Publications & CD-ROMs

Graphs & Maps

Multimedia

Video

Databases & Software

AquaCrop

CropWat

ClimWat

ETo Calculator

Wastewater

Projects

Hot Issues

QUICKLINK TO ▾

Databases & Software

Publications & CD-ROMs

Educational Material

Our Regional Offices

IPTRID


Databases

CLIMWAT 2.0 for CROPWAT

CLIMWAT is a climatic database to be used in combination with the computer program CROPWAT, and allows the calculation of crop water requirements, irrigation supply and irrigation scheduling for various crops for a range of climatological stations worldwide.

CLIMWAT 2.0 for CROPWAT is a joint publication of the **Water Development and Management Unit** and the **Climate Change and Bioenergy Unit** of FAO.

CLIMWAT 2.0 offers observed agroclimatic data of over 5000 stations worldwide distributed as shown below.



Location of stations included in CLIMWAT 2.0.

SEARCH

RELATED LINKS

Climate Change and Bioenergy Unit

Revised FAO Methodology for Crop Water Requirements

DOWNLOADS

ClimWat 2.0 for Cropwat

CONTACT US





Des Données d'Evapotranspiration

http://www.fao.org/nr/water/inforest_databases_cropwat.html

▶ text only version ▶ print friendly helping to build a world without hunger

NATURAL RESOURCES AND ENVIRONMENT DEPARTMENT



FAO WATER

FrontPage

Topics

Information Resources

Publications & CD-ROMs

Graphs & Maps

Multimedia

Video

Databases & Software

AquaCrop

CropWat

ClimWat

ETo Calculator

Wastewater

Projects

Hot Issues

QUICKLINK TO ↴

Databases & Software

Publications & CD-ROMs

Educational Material

Our Regional Offices

IPTRID

Software

CropWat

CROPWAT is a decision support tool developed by the Land and Water Development Division of FAO.

CROPWAT 8.0 for Windows is a computer program for the calculation of crop water requirements and irrigation requirements based on soil, climate and crop data. In addition, the program allows the development of irrigation schedules for different management conditions and the calculation of scheme water supply for varying crop patterns. CROPWAT 8.0 can also be used to evaluate farmers' irrigation practices and to estimate crop performance under both rainfed and irrigated conditions.

All calculation procedures used in CROPWAT 8.0 are based on the two FAO publications of the Irrigation and Drainage Series, namely, No. 56 "Crop Evapotranspiration - Guidelines for computing crop water requirements" and No. 33 titled "Yield response to water".

As a starting point, and only to be used when local data are not available, CROPWAT 8.0 includes standard crop and soil data. When local data are available, these data files can be easily modified or new ones can be created. Likewise, if local climatic data are not available, these can be obtained for over 5,000 stations worldwide from CLIMWAT, the associated climatic database. The development of irrigation schedules in CROPWAT 8.0 is based on the FAO Irrigation Scheduling Methodology.



SEARCH

RELATED LINKS

Revised methodology for crop water requirements

Topics: Irrigation

Topics: Water Productivity

Publications: Irrigation

Publications: Productivity

DOWNLOADS

CropWat 8.0

Example of CropWat 8.0 use

CONTACT US



Pratiques d'irrigation et les rendements agricoles

http://www.fao.org/ag/agl/fertistat/fst_fubc_en.asp





Volumes de production et le commerce

<http://faostat.fao.org/default.aspx>

Google FAO: Search result FAO - Water Development FAO - Water Development FAOSTAT

faostat.fao.org/default.aspx

FOOD AND AGRICULTURE

FAOSTAT

Home Production Trade Food Supply Food Balance Sheets Food Security Prices Resources Forestry Fisheries Metadata Support/FAC

logon want to register? FAOSTAT videos CountrySTAT

FAOSTAT provides time-series and cross sectional data relating to food and agriculture for some 200 countries.

The national version of FAOSTAT, **CountrySTAT**, is being developed and implemented in a number of target countries, primarily in sub-saharan Africa. It will offer a two-way data exchange facility between countries and FAO as well as a facility to store data at the national and sub-national levels.

FAOSTAT User Dialogue 2010

One of the priorities of the Statistics Division is to involve the producers and custodians of statistics at FAO in a stronger communication with our users, and to gain a better understanding of their needs. Since July 2010 the free FAO data access policy has been implemented. The number of registered users has boomed since then from 400 to over 10 000 as of today.

We are seeking contact with registered users of FAOSTAT to open a qualitative feedback channel. We are looking for your views and experiences in using FAO's statistics as well as specific, technical suggestions. We would very much appreciate receiving your advice. To see the issues please click [here](#).

Latest news

The following subject domains have been updated:

- [ForesSTAT \(August 2011\)](#)
- [PopSTAT \(August 2011\)](#)
- [ResourceSTAT-Land \(July 2011\)](#)
- [Forestry Trade Flows \(July 2011\)](#)
- [ResourceSTAT-Fertilizers \(July 2011\)](#)
- [ProdSTAT \(May 2011\)](#)
- [ResourceSTAT-Pesticides Consumption \(March 2011\)](#)
- [TradeSTAT-Detailed Trade Matrix \(February 2011\)](#)
- [ResourceSTAT-Machinery \(December 2010\)](#)
- [TradeSTAT-Detailed trade data \(November 2010\)](#)



Water Footprint Network

<http://www.waterfootprint.org>





Autres références utiles

Allen, R.G., Pereira, L.S., Raes, D. and Smith, M. (1998) Crop evapotranspiration - Guidelines for computing crop water requirements - FAO Irrigation and drainage paper 56. Food and Agriculture Organization. Rome. Available from: <http://www.fao.org/docrep/X0490E/X0490E00.htm>

Chapagain, A.K. and Hoekstra, A.Y. (2004) Water footprints of nations. Value of Water Research Report Series No. 16, UNESCO-IHE, Delft, the Netherlands. Available from:
<http://www.waterfootprint.org/?page=files/Research%20data>

Mekonnen, M.M. and Hoekstra, A.Y. (2011) The green, blue and grey water footprint of crops and derived crop products, Hydrology and Earth System Sciences, 15(5): 1577-1600. Available from:

<http://www.waterfootprint.org/?page=files/Publications>

Chapagain, A.K. and Hoekstra, A.Y. (2003) 'Virtual water flows between nations in relation to trade in livestock and livestock products' ,Value of Water Research Report Series No.13, UNESCO-IHE.

<http://www.waterfootprint.org/?page=files/Publications>

FAO (2003) Technical Conversion Factors for Agricultural Commodities. Food and Agriculture Organization of the United Nations, Rome. Online available: www.fao.org/WAICENT/FAOINFO/ECONOMIC/ESS/pdf/tcf.pdf

Hoekstra, A.Y. and Chapagain, A.K. (2008) Globalization of water: Sharing the planet's freshwater resources. Blackwell Publishing. Oxford, UK.

Monfreda, C., Ramankutty, N. and Foley, J.A. (2008) Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, Global Biogeochemical Cycles, Vol. 22, GB1022, doi:10.1029/2007GB002947. Available from:
<http://www.geog.mcgill.ca/~nramankutty/Datasets/Datasets.html>



4: Utilisant CROPWAT





$$WF = \frac{\text{Crop water use (m}^3/\text{ha})}{\text{Crop yield (ton/ha)}}$$

CROPWAT
(FAO, 2009a)


FAOSTAT
(FAO, 2009b)

CROPWAT estimation methods

1. CWR option
 - Optimal conditions
 - Without considering soil water balance
 - Ten-day time step effective rainfall
2. Irrigation schedule option
 - Non-optimal conditions
 - Considering soil water balance
 - Daily time step soil water balance




1. CWR option





1. CWR option

Reference (ET_0), crop evapotranspiration under standard (ET_C) and non-standard conditions ($ET_{C\ adj}$)



Reference crop evapotranspiration

Evapotranspiration rate from a hypothetical grass reference surface, not short of water

Crop evapotranspiration under standard conditions

Evapotranspiration from disease-free, well fertilized crops, grown in large fields, under optimum soil water conditions and achieving full production under the given climatic conditions


Allen et al. (1998)

Crop coefficient

Characteristics: Crop height, albedo (reflectance), canopy resistance, evaporation from soil.




1. CWR option – Input





Climate data – CLIMWAT 2.0 database (FAO, 2006)


Observed agroclimatic data of over 5000 stations worldwide (tries to cover 1971-2000)





Climate data – **New LocClim** 1.10 database (FAO, 2005)

Interpolates where no observations are available





Climate data – Local Data can be added by hand to CROPWAT





CLIMWAT database

.CLI

GRONINGEN-AP-EELDE.cli - Notepad		
File	Edit	Format View Help
"GRONINGEN-AP-EELDE", "", " 4 ", 0, 0, 0, 0, 0, 0, 0, 0		
0.21	66.60	59.50
0.39	45.20	41.93
0.86	57.50	52.21
1.58	48.20	44.48
2.43	57.80	52.45
2.82	68.80	61.23
2.75	76.30	66.99
2.53	66.40	59.35
1.59	70.60	62.63
0.83	68.70	61.15
0.44	77.10	67.59
0.22	75.50	66.38

*Penman-Monteith method

**USDA Soil Conservation Service formula

ET_0^* (mm/day)	Monthly rainfall (mm/month)	Monthly effective rainfall** (mm/month)
----------------------	-----------------------------------	---

.PEN

GRONINGEN-AP-EELDE.pen - Notepad						
File	Edit	Format	View	Help		
"Location 7", "GRONINGEN-AP-EELDE", 4, 53.13, "N.L.", 6.58, " 01 "						
3.7	-1.5	95.4	457.9	0.45	2.06	0.21
4.5	-1.4	91.9	423.4	2.05	4.60	0.39
7.9	0.5	86.7	449.3	2.85	7.93	0.86
11.8	2.5	81.7	397.4	5.01	13.22	1.58
16.6	6.3	79.1	354.2	6.22	17.06	2.43
19.5	9.3	80.6	345.6	6.11	17.95	2.82
20.7	10.9	83.0	354.2	5.71	16.92	2.75
21.2	10.8	82.5	319.7	5.83	15.05	2.53
18.2	8.7	87.2	337.0	4.06	10.18	1.59
13.7	5.9	90.8	354.2	2.45	5.78	0.83
8.3	2.3	92.5	440.6	1.06	2.77	0.44
4.9	-0.2	95.7	449.3	0.13	1.53	0.22

Mean daily maximum temp. (°C)	Mean daily minimum temp. (°C)	Mean relative humidity (%)	Mean wind speed (km/day)	Mean sunshine hours per day	Mean solar radiation (MJ/m ² /day)	ET_0^* (mm/day)
-------------------------------	-------------------------------	----------------------------	--------------------------	-----------------------------	---	-------------------





ET_0 Penman-Monteith

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ETo

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

ETo file Rain file Crop file Soil file Pla

CROPWAT options

Climate / ETo Rainfall Non-rice crop scheduling Rice scheduling Land Preparation (rice)

Data settings

ETo Penman-Monteith ETo Penman calculated from climatic data

Temperature Minimum / maximum temperatures

Changes to these settings only effect NEW data

Units

Humidity Relative humidity in %

Windspeed Kilometers per day

Sunshine Hours sunshine

ETo mm per day

Save as default Reset to FAO defaults OK Cancel Help





Effective rainfall estimation method

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ETo

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

CROPWAT options

- Climate / ETo
- Rainfall
- Non-rice crop scheduling
- Rice scheduling
- Land Preparation (rice)

Effective rainfall method for CWR calculations

Fixed Percentage: 80 %

Dependable rain (FAO/AGLW formula)
 $P_{eff} = 0.6 \times P - 10 \frac{P}{3}$ for $P_{month} \leq 70 \frac{P}{3}$ mm
 $P_{eff} = 0.8 \times P - 24 \frac{P}{3}$ for $P_{month} > 70 \frac{P}{3}$ mm

Empirical formula
 $P_{eff} = 0.5 \times P + -5 \frac{P}{3}$ for $P \leq 50 \frac{P}{3}$ mm
 $P_{eff} = 0.7 \times P + 20 \frac{P}{3}$ for $P > 50 \frac{P}{3}$ mm

USDA soil conservation service
 $P_{eff} = (P * (125 - 0.2 \times P)) / 125$ for $P \leq 250 \frac{P}{3}$ mm
 $P_{eff} = 125 \frac{P}{3} + 0.1 \times P$ for $P > 250 \frac{P}{3}$ mm

Rainfall not considered in irrigation calculations (effective rainfall = 0)

Note: in red are correction factors that CROPWAT applies to adjust formulas in the case of decade and daily rainfall data (for effective rainfall calculations daily data are aggregated per decade)

Save as default Reset to FAO defaults OK Cancel Help

ETo file	Rain file	Crop file	Soil file	Pla
----------	-----------	-----------	-----------	-----

→ Climate/ETo

→ Rain



Crop parameters


- Crop coefficients (K_c)
 - Global → CROPWAT (FAO, 2009a)
 - Regional → Table 12 in Allen et al. (1998)
 - Climate region → Chapagain and Hoekstra (2004) (App. VI)
 - Local → Agricultural research stations, farmers

- Crop growing period
(planting/green up date)
 - Global → CROPWAT (FAO, 2009a)
 - Regional → Table 11 in Allen et al. (1998)
 - Climate region → Chapagain and Hoekstra (2004) (App. VI)
 - Local → Agricultural research stations, farmers





Crop coefficient curve




Allen et al. (1998)





Crop growth stages for different crops

Planting date



Allen et al. (1998)





1. CWR option - input

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ET₀

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

E_{To} file Rain file Crop file Soil file Planting date

The screenshot shows the CROPWAT software interface. The title bar reads "CROPWAT - Session: untitled". The menu bar includes File, Edit, Calculations, Charts, Settings, Window, Language, and Help. The toolbar below the menu bar contains icons for New, Open, Save, Close, Print, Chart, and Options. On the left, a vertical sidebar lists several options with corresponding icons: Climate/ET₀, Rain, Crop, Soil, CWR (which is highlighted with a yellow background), Schedule, Crop Pattern, and Scheme. Below this sidebar is a row of six buttons labeled "E_{To} file", "Rain file", "Crop file", "Soil file", "Planting date", and an empty button. A large empty area to the right of the sidebar is where data would typically be entered or displayed.





1. CWR option - output

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ET₀

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

E_{To} file Rain file Crop file Soil file Planting date





1. CWR option - output

Crop Water Requirements

ET ₀ station		Rain station		CWR		Crop	
GRONINGEN-AP-EELD		GRONINGEN-AP-EELD				Sugarbeet	
						Planting date	
						10/04	
Month	Decade	Stage	K _c	ET _c	ET _c	Eff rain	Irr. Req.
			coeff	mm/day	mm/dec	mm/dec	mm/dec
Apr	1	Init	0.35	0.52	0.5	1.5	0.5
Apr	2	Init	0.35	0.62	6.2	14.1	0.0
Apr	3	Init	0.35	0.72	7.2	15.2	0.0
May	1	Init	0.35	0.83	8.3	16.6	0.0
May	2	Init	0.35	0.93	9.3	17.5	0.0
May	3	Deve	0.36	1.00	11.0	18.5	0.0
Jun	1	Deve	0.51	1.51	15.1	19.5	0.0
Jun	2	Deve	0.73	2.25	22.5	20.5	2.0
Jun	3	Deve	0.95	2.90	29.0	21.1	7.9
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6
Aug	1	Mid	1.23	3.50	35.0	20.3	14.7
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8
Aug	3	Late	1.22	2.97	32.7	19.8	12.9
Sep	1	Late	1.11	2.33	23.3	20.6	2.7
Sep	2	Late	0.98	1.73	17.3	21.1	0.0
Sep	3	Late	0.86	1.27	12.7	20.8	0.0
Oct	1	Late	0.76	0.91	5.5	12.2	0.0
					381.2	345.6	100.1

→ 



1. CWR option - output

Crop Water Requirements

ET ₀ station		GRONINGEN-AP-EELD				Crop		Sugarbeet
Rain station		GRONINGEN-AP-EELD				Planting date		10/04
				CWR				
Month	Decade	Stage	K _c	ET _c	ET _c	Eff rain	Irr. Req.	
			coeff	mm/day	mm/dec	mm/dec	mm/dec	
Apr	1	Init	0.35	0.52	0.5	1.5	0.5	
Apr	2	Init	0.35	0.62	6.2	14.1	0.0	
Apr	3	Init	0.35	0.72	7.2	15.2	0.0	
May	1	Init	0.35	0.83	8.3	16.6	0.0	
May	2	Init	0.35	0.93	9.3	17.5	0.0	
May	3	Deve	0.36	1.00	11.0	18.5	0.0	
Jun	1	Deve	0.51	Copy table	Data only		0.0	
Jun	2	Deve	0.73	2.25	Data and Headers		2.0	
Jun	3	Deve	0.95	2.90	29.0	21.1	7.9	
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1	
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8	
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6	
Aug	1	Mid	1.23	3.50	35.0	20.3	14.7	
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8	
Aug	3	Late	1.22	2.97	32.7	19.8	12.9	
Sep	1	Late	1.11	2.33	23.3	20.6	2.7	
Sep	2	Late	0.98	1.73	17.3	21.1	0.0	
Sep	3	Late	0.86	1.27	12.7	20.8	0.0	
Oct	1	Late	0.76	0.91	5.5	12.2	0.0	
					381.2	345.6	100.1	





1. CWR option - output

$$ET = ET_c$$

$$ET_g = \min (ET_c, P_{eff})$$

$ET_b = \max (0, ET_c - P_{eff}) * \text{Irr. Fraction}$ (ie the volume of irrigation in reality)


CWR

Month	Decade	Stage	Kc	ETc	ETc	Eff rain	Irr. Req.	ET	ETg	ETb
			coeff	mm/day	mm/dec	mm/dec	mm/dec	mm/dec	mm/dec	mm/dec
Apr	1	Init	0.35	0.52	0.5	1.5	0.5	0.5	0.5	0
Apr	2	Init	0.35	0.62	6.2	14.1	0	6.2	6.2	0
Apr	3	Init	0.35	0.72	7.2	15.2	0	7.2	7.2	0
May	1	Init	0.35	0.83	8.3	16.6	0	8.3	8.3	0
May	2	Init	0.35	0.93	9.3	17.5	0	9.3	9.3	0
May	3	Deve	0.36	1	11	18.5	0	11	11	0
Jun	1	Deve	0.51	1.51	15.1	19.5	0	15.1	15.1	0
Jun	2	Deve	0.73	2.25	22.5	20.5	2	22.5	20.5	2
Jun	3	Deve	0.95	2.9	29	21.1	7.9	29	21.1	7.9
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1	35.2	22.1	13.1
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8	36.7	22.9	13.8
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6	39.5	21.9	17.6
Aug	1	Mid	1.23	3.5	35	20.3	14.7	35	20.3	14.7
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8	34.1	19.3	14.8
Aug	3	Late	1.22	2.97	32.7	19.8	12.9	32.7	19.8	12.9
Sep	1	Late	1.11	2.33	23.3	20.6	2.7	23.3	20.6	2.7
Sep	2	Late	0.98	1.73	17.3	21.1	0	17.3	17.3	0
Sep	3	Late	0.86	1.27	12.7	20.8	0	12.7	12.7	0
Oct	1	Late	0.76	0.91	5.5	12.2	0	5.5	5.5	0
Total					381	346	100	381	282	100






2. Irrigation schedule option






2. Irrigation schedule option





2. Irrigation schedule option input





Soil data (FAO, 2009a)

Soil - C:\ProgramData\CROPWAT\data\soils\FAO\MEDIUM.SOI

Soil name	Medium (loam)
General soil data	
Total available soil moisture (FC - WP)	290.0 mm/meter
Maximum rain infiltration rate	40 mm/day
Maximum rooting depth	900 centimeters
Initial soil moisture depletion (as % TAM)	0 %
Initial available soil moisture	290.0 mm/meter





Soil data (FAO, 2009a)

Field Capacity (FC)

- The amount of moisture that remains in the soil after the excess water has drained away.
- FC varies with soil type

Wilting Point (WP)

- The minimum amount of soil moisture required for a plant to remain upright.
- If soil contains less moisture than the wilting point for that plant, it will wilt, but can recover.

Total Available Water

- Water held in the soil available to plants
- Difference between the water content at field capacity and wilting point

Readily Available Water

- Fraction of TAW that a crop can extract from the root zone without suffering water stress



Actual evapotranspiration


Reference
evapotranspiration
 ET_o



Crop water requirement
 CWR
 $CWR = ET_o \times K_c$




Actual
evapotranspiration ET_a
 $ET_a = CWR \times K_s$




Soil

Soil water balance




Soil water balance






2. Irrigation schedule option – input





2. Irrigation schedule option – output





2. Irrigation schedule option - output

Crop irrigation schedule

ET ₀ station	GRONINGEN-AP-EELD	Crop	Sugarbeet	Planting date	10/04	Yield red.	
Rain station	GRONINGEN-AP-EELD	Soil	Medium (loam)	Harvest date	06/10		0.0 %

Table format

Irrigation schedule Daily soil moisture balance

Timing: Irrigate at critical depletion
Application: Refill soil to field capacity
Field eff. 70 %

Date	Day	Stage	Rain	K _s	Eta	Depl	Net Irr	Deficit	Loss	Gr. Irr
			mm	fract.	mm/day	%	mm	mm	mm	mm
10 Apr	1	Init	0.0	1.00	0.5	1	0.0	0.5	0.0	0.0
11 Apr	2	Init	0.0	1.00	0.6	1	0.0	1.1	0.0	0.0
12 Apr	3	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
13 Apr	4	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
14 Apr	5	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0
15 Apr	6	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
16 Apr	7	Init	0.0	1.00	0.6	2	0.0	2.5	0.0	0.0
17 Apr	8	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
18 Apr	9	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0

Totals

Total gross irrigation	0.0	mm	Total rainfall	383.0	mm
Total net irrigation	0.0	mm	Effective rainfall	319.5	mm
Total irrigation losses	0.0	mm	Total rain loss	63.4	mm
Actual water use by crop	380.3	mm	Moist deficit at harvest	60.8	mm
Potential water use by crop	380.3	mm	Actual irrigation requirement	60.8	mm
Efficiency irrigation schedule	-	%	Efficiency rain	83.4	%
Deficiency irrigation schedule	0.0	%			

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------

→ Schedule



2. Irrigation schedule option - output

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Crop irrigation schedule

Climate/ETo ETo station GRONINGEN-AP-EELD Crop Sugarbeet Planting date 10/04 Yield red.
Rain station GRONINGEN-AP-EELD Soil Medium (loam) Harvest date 06/10 0.0 %

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

Table format

Irrigation schedule
 Daily soil moisture balance

Timing: Irrigate at critical depletion
Application: Refill soil to field capacity
Field eff. 70 %

Date	Day	Stage	Rain	Ks	Eta	Depl	Net Irr	Deficit	Loss	Gr. Irr
10 Apr	1	Init	0.0	1.00	0.5	1	0.0	0.5	0.0	0.0
11 Apr	2	Init	0.0	1.00	0.6	1	0.0	1.1	0.0	0.0
12 Apr	3	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
13 Apr	4	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
14 Apr	5	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0
15 Apr	6	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
16 Apr	7	Init	0.0	1.00	0.6	2	0.0	2.5	0.0	0.0
17 Apr	8	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
18 Apr	9	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0

Totals

Total gross irrigation	0.0 mm	Total rainfall	383.0 mm
Total net irrigation	0.0 mm	Effective rainfall	319.5 mm
Total irrigation losses	0.0 mm	Total rain loss	63.4 mm
Actual water use by crop	380.3 mm	Moist deficit at harvest	60.8 mm
Potential water use by crop	380.3 mm	Actual irrigation requirement	60.8 mm
Efficiency irrigation schedule	- %	Efficiency rain	83.4 %
Deficiency irrigation schedule	0.0 %		

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------





2. Irrigation schedule option - output

Rainfed conditions

$$ET_{green} (irr = 0) = ET_{tot} (irr = 0)$$

$$ET_{blue} (irr = 0) = 0$$

Irrigated conditions

$$ET_{green} (irr = 1) = ET_{green} (irr = 0)$$

$$ET_{blue} (irr = 1) = ET_{tot} (irr = 1) - ET_{green} (irr = 0)$$

Rainfed scenario (*irr = 0*)

CROPWAT options

Non-rice crop scheduling |

Scheduling criteria for non-rice crops

Irrigation timing

No irrigation (rainfed)

No irrigation (only rainfall)

Irrigation application

Refill soil to field capacity

Irrigation efficiency

Irrigation efficiency: 70 %

Save as default Reset to FAO defaults OK Cancel Help

Irrigated scenario (*irr = 1*)

CROPWAT options

Non-rice crop scheduling |

Scheduling criteria for non-rice crops

Irrigation timing

Irrigate at critical depletion

Irrigation at 100 % critical depletion

Irrigation application

Refill soil to field capacity

Irrigation efficiency

Irrigation efficiency: 70 %

Save as default Reset to FAO defaults OK Cancel Help




Calculating the process water footprint of growing a crop: An example for sugar beet in Northern Netherlands





Climate data - CLIMWAT database (FAO, 2006)


Observed agroclimatic data of over 5000 stations worldwide (tries to cover 1971-2000)



Green and blue water footprint of a crop



Crop distribution maps




Sugar beet harvested area in the Netherlands
(unit: proportion of grid cell area) Source: Monfreda et al. (2008)





Crop distribution maps – Climate data




Climate stations (dots) and sugar beet harvested area in the Netherlands
(unit: proportion of grid cell area) Source: FAO (2006) Monfreda et al. (2008)






Crop distribution maps – Climate data



Climate stations (dots) and sugar beet harvested area in the Netherlands
(unit: proportion of grid cell area) Source: FAO (2006) Monfreda et al. (2008)



Crop distribution maps – Climate data



Groningen Ap Eelde

Climate stations (dots) and sugar beet harvested area in the Netherlands
(unit: proportion of grid cell area) Source: FAO (2006) Monfreda et al. (2008)





Climate data - CLIMWAT

.CLI

GRONINGEN-AP-EELDE.cli - Notepad		
File	Edit	Format
View	Help	
"GRONINGEN-AP-EELDE", "", " 4", 0, 0, 0, 0, 0, 0, 0		
0.21	66.60	59.50
0.39	45.20	41.93
0.86	57.50	52.21
1.58	48.20	44.48
2.43	57.80	52.45
2.82	68.80	61.23
2.75	76.30	66.99
2.53	66.40	59.35
1.59	70.60	62.63
0.83	68.70	61.15
0.44	77.10	67.59
0.22	75.50	66.38

*Penman-Monteith method

**USDA Soil Conservation Service formula

ET ₀ [*] (mm/day)	Monthly rainfall (mm/month)	Monthly effective rainfall** (mm/month)
--	-----------------------------------	---

.PEN

GRONINGEN-AP-EELDE.pen - Notepad						
File	Edit	Format	View	Help		
"Location 7", "GRONINGEN-AP-EELDE", 4, 53.13, "N.L.", 6.58, " 01"						
3.7	-1.5	95.4	457.9	0.45	2.06	0.21
4.5	-1.4	91.9	423.4	2.05	4.60	0.39
7.9	0.5	86.7	449.3	2.85	7.93	0.86
11.8	2.5	81.7	397.4	5.01	13.22	1.58
16.6	6.3	79.1	354.2	6.22	17.06	2.43
19.5	9.3	80.6	345.6	6.11	17.95	2.82
20.7	10.9	83.0	354.2	5.71	16.92	2.75
21.2	10.8	82.5	319.7	5.83	15.05	2.53
18.2	8.7	87.2	337.0	4.06	10.18	1.59
13.7	5.9	90.8	354.2	2.45	5.78	0.83
8.3	2.3	92.5	440.6	1.06	2.77	0.44
4.9	-0.2	95.7	449.3	0.13	1.53	0.22

Mean daily maximum temp. (°C)	Mean daily minimum temp. (°C)	Mean relative humidity (%)	Mean wind speed (km/day)	Mean sunshine hours per day	Mean solar radiation (MJ/m ² /day)	ET (mm)	Rain
-------------------------------	-------------------------------	----------------------------	--------------------------	-----------------------------	---	---------	------





Crop parameters for sugar beet


Source	Level	Kc ini	Kc mid	Kc end	Initial stage	Dev stage	Mid stage	Late stage	Planting/Green up date	Rooting depth (m)	Critical depletion level (p)	Yield response factor (Ky)
Chapagain and Hoekstra (2004)	Per Climate Region	0.35	1.20	0.70	50	40	50	40	1 April			
Allen et al. (1998)	Per region	0.35	1.20	0.70	50	40	50	40	April			
CROPWAT (FAO, 2009a)	Global	0.35	1.20	0.70	25	35	50	50		0.3-1	0.5-0.6-0.6	0.5-0.8-1.2-1-1.1
IRS (2009)*	Local								10 April			

*Institute of Sugar Beet Research in the Netherlands






Crop parameters for sugar beet





Crop parameters for sugar beet





Crop parameters for sugarbeet

```
SUGARBET.CRO - Notepad
File Edit Format View Help
CROPWAT 8.0 Crop data
Sugarbeet
50 40 50 40
 0.35 1.20 0.70
 0.30 1.00
 0.50 0.60 0.60
 0.50 0.80 1.20 1.00 1.10
 0.70
```

- Crop length of the growing season
- Crop coefficients (Kc init, Kc mid, Kc end)
- Rooting depth (m)
- Critical depletion level (P)
- Yield response factor (Ky)
- Height (provide if available)





1. CWR option - input

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ET₀

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

ETo file Rain file Crop file Soil file Planting date

The screenshot shows the CROPWAT software window titled "CROPWAT - Session: untitled". The menu bar includes File, Edit, Calculations, Charts, Settings, Window, Language, and Help. The toolbar below the menu has buttons for New, Open, Save, Close, Print, Chart, and Options. On the left, a vertical sidebar lists several options with icons: Climate/ET₀, Rain, Crop, Soil, CWR (selected), Schedule, Crop Pattern, and Scheme. Below this sidebar is a row of six input fields labeled ETo file, Rain file, Crop file, Soil file, Planting date, and an empty field. The "CWR" option is highlighted with a blue selection bar.





1. CWR option - output

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Climate/ET₀

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

ETo file Rain file Crop file Soil file Planting date

The screenshot shows the CROPWAT software interface. The title bar reads "CROPWAT - Session: untitled". The menu bar includes File, Edit, Calculations, Charts, Settings, Window, Language, and Help. Below the menu is a toolbar with icons for New, Open, Save, Close, Print, Chart, and Options. On the left, a vertical sidebar lists several options: Climate/ET₀, Rain, Crop, Soil, CWR (which is highlighted with a grey arrow), Schedule, Crop Pattern, and Scheme. At the bottom, there are six input fields labeled ETo file, Rain file, Crop file, Soil file, Planting date, and an empty field. The "CWR" option is currently active.





1. CWR option - output

Crop Water Requirements

ET ₀ station			Crop				
GRONINGEN-AP-EELD			Sugarbeet				
Rain station			Planting date				
GRONINGEN-AP-EELD			10/04				
Month	Decade	Stage	K _c coeff	ET _c mm/day	ET _c mm/dec	Eff rain mm/dec	Irr. Req. mm/dec
Apr	1	Init	0.35	0.52	0.5	1.5	0.5
Apr	2	Init	0.35	0.62	6.2	14.1	0.0
Apr	3	Init	0.35	0.72	7.2	15.2	0.0
May	1	Init	0.35	0.83	8.3	16.6	0.0
May	2	Init	0.35	0.93	9.3	17.5	0.0
May	3	Deve	0.36	1.00	11.0	18.5	0.0
Jun	1	Deve	0.51	1.51	15.1	19.5	0.0
Jun	2	Deve	0.73	2.25	22.5	20.5	2.0
Jun	3	Deve	0.95	2.90	29.0	21.1	7.9
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6
Aug	1	Mid	1.23	3.50	35.0	20.3	14.7
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8
Aug	3	Late	1.22	2.97	32.7	19.8	12.9
Sep	1	Late	1.11	2.33	23.3	20.6	2.7
Sep	2	Late	0.98	1.73	17.3	21.1	0.0
Sep	3	Late	0.86	1.27	12.7	20.8	0.0
Oct	1	Late	0.76	0.91	5.5	12.2	0.0
				381.2	345.6	100.1	
				IR = ET _c -P _{eff}			





1. CWR option - output

Crop Water Requirements

ETO station: GRONINGEN-AP-EELD Crop: Sugarbeet

Rain station: GRONINGEN-AP-EELD Planting date: 10/04

Month	Decade	Stage	Kc coeff	ETc mm/day	ETc mm/dec	Eff rain mm/dec	Irr. Req. mm/dec
Apr	1	Init	0.35	0.52	0.5	1.5	0.5
Apr	2	Init	0.35	0.62	6.2	14.1	0.0
Apr	3	Init	0.35	0.72	7.2	15.2	0.0
May	1	Init	0.35	0.83	8.3	16.6	0.0
May	2	Init	0.35	0.93	9.3	17.5	0.0
May	3	Deve	0.36	1.00	11.0	18.5	0.0
Jun	1	Deve	0.51	Copy table ▶	Data only		0.0
Jun	2	Deve	0.73	2.25	Data and Headers		2.0
Jun	3	Deve	0.95	2.90	29.0	21.1	7.9
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6
Aug	1	Mid	1.23	3.50	35.0	20.3	14.7
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8
Aug	3	Late	1.22	2.97	32.7	19.8	12.9
Sep	1	Late	1.11	2.33	23.3	20.6	2.7
Sep	2	Late	0.98	1.73	17.3	21.1	0.0
Sep	3	Late	0.86	1.27	12.7	20.8	0.0
Oct	1	Late	0.76	0.91	5.5	12.2	0.0
					381.2	345.6	100.1

▶

CWR



1. CWR option - output

$$ET = ET_c$$

$$ET_g = \min (ET_c, P_{\text{eff}})$$

$$ET_b = \max (0, ET_c - P_{\text{eff}}) * \text{Irr. fraction}$$

Growing period

Month	Decade	Stage	Kc coeff	ETc mm/day	ETc mm/dec	Eff rain mm/dec	Irr. Req. mm/dec	ET mm/dec	ETg mm/dec	ETb mm/dec
Apr	1	Init	0.35	0.52	0.5	1.5	0.5	0.5	0.5	0
Apr	2	Init	0.35	0.62	6.2	14.1	0	6.2	6.2	0
Apr	3	Init	0.35	0.72	7.2	15.2	0	7.2	7.2	0
May	1	Init	0.35	0.83	8.3	16.6	0	8.3	8.3	0
May	2	Init	0.35	0.93	9.3	17.5	0	9.3	9.3	0
May	3	Deve	0.36	1	11	18.5	0	11	11	0
Jun	1	Deve	0.51	1.51	15.1	19.5	0	15.1	15.1	0
Jun	2	Deve	0.73	2.25	22.5	20.5	2	22.5	20.5	2
Jun	3	Deve	0.95	2.9	29	21.1	7.9	29	21.1	7.9
Jul	1	Mid	1.17	3.52	35.2	22.1	13.1	35.2	22.1	13.1
Jul	2	Mid	1.23	3.67	36.7	22.9	13.8	36.7	22.9	13.8
Jul	3	Mid	1.23	3.59	39.5	21.9	17.6	39.5	21.9	17.6
Aug	1	Mid	1.23	3.5	35	20.3	14.7	35	20.3	14.7
Aug	2	Mid	1.23	3.41	34.1	19.3	14.8	34.1	19.3	14.8
Aug	3	Late	1.22	2.97	32.7	19.8	12.9	32.7	19.8	12.9
Sep	1	Late	1.11	2.33	23.3	20.6	2.7	23.3	20.6	2.7
Sep	2	Late	0.98	1.73	17.3	21.1	0	17.3	17.3	0
Sep	3	Late	0.86	1.27	12.7	20.8	0	12.7	12.7	0
Oct	1	Late	0.76	0.91	5.5	12.2	0	5.5	5.5	0
Total					381	346	100	381	282	100





1. CWR option

Water footprint estimation


ET_g	ET_b	ET	CWU_g	CWU_b	CWU	Y^*	$WF_{proc,green}$	$WF_{proc,blue}$	WF_{proc}
mm/period			m^3/ha			ton/ha	m^3/ton		
282	100	381	2816	995	3811	59	48	17	65

*IRS (1999-2006) for Northern Netherlands






2. Irrigation schedule option – input





2. Irrigation schedule option – output





2. Irrigation schedule option - output

Crop irrigation schedule

ET ₀ station	GRONINGEN-AP-EELD	Crop	Sugarbeet	Planting date	10/04	Yield red.	
Rain station	GRONINGEN-AP-EELD	Soil	Medium (loam)	Harvest date	06/10		0.0 %

Table format:

<input type="radio"/> Irrigation schedule	Timing: Irrigate at critical depletion
<input checked="" type="radio"/> Daily soil moisture balance	Application: Refill soil to field capacity
	Field eff. 70 %

Date Day Stage Rain Ks Eta Depl Net Irr Deficit Loss Gr. Irr

mm fract. mm/day % mm mm mm mm

10 Apr	1	Init	0.0	1.00	0.5	1	0.0	0.5	0.0	0.0
11 Apr	2	Init	0.0	1.00	0.6	1	0.0	1.1	0.0	0.0
12 Apr	3	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
13 Apr	4	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
14 Apr	5	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0
15 Apr	6	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
16 Apr	7	Init	0.0	1.00	0.6	2	0.0	2.5	0.0	0.0
17 Apr	8	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
18 Apr	9	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0

Totals

Total gross irrigation	0.0	mm	Total rainfall	383.0	mm
Total net irrigation	0.0	mm	Effective rainfall	319.5	mm
Total irrigation losses	0.0	mm	Total rain loss	63.4	mm
Actual water use by crop	380.3	mm	Moist deficit at harvest	60.8	mm
Potential water use by crop	380.3	mm	Actual irrigation requirement	60.8	mm
Efficiency irrigation schedule	-	%	Efficiency rain	83.4	%
Deficiency irrigation schedule	0.0	%			

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------





2. Irrigation schedule option - output

CROPWAT - Session: untitled

File Edit Calculations Charts Settings Window Language Help

New Open Save Close Print Chart Options

Crop irrigation schedule

Climate/ETo ETo station GRONINGEN-AP-EELD Crop Sugarbeet Planting date 10/04 Yield red.
Rain station GRONINGEN-AP-EELD Soil Medium (loam) Harvest date 06/10 0.0 %

Rain

Crop

Soil

CWR

Schedule

Crop Pattern

Scheme

Table format

Irrigation schedule Daily soil moisture balance

Timing: Irrigate at critical depletion
Application: Refill soil to field capacity
Field eff. 70 %

Date	Day	Stage	Rain	Ks	Eta	Depl	Net Irr	Deficit	Loss	Gr. Irr
			mm	fract.	mm/day	%	mm	mm	mm	mm
10 Apr	1	Init	0.0	1.00	0.5	1	0.0	0.5	0.0	0.0
11 Apr	2	Init	0.0	1.00	0.6	1	0.0	1.1	0.0	0.0
12 Apr	3	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
13 Apr	4	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
14 Apr	5	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0
15 Apr	6	Init	0.0	1.00	0.6	2	0.0	1.8	0.0	0.0
16 Apr	7	Init	0.0	1.00	0.6	2	0.0	2.5	0.0	0.0
17 Apr	8	Init	7.6	1.00	0.6	1	0.0	0.6	0.0	0.0
18 Apr	9	Init	0.0	1.00	0.6	1	0.0	1.2	0.0	0.0

Totals

Total gross irrigation	0.0 mm	Total rainfall	383.0 mm
Total net irrigation	0.0 mm	Effective rainfall	319.5 mm
Total irrigation losses	0.0 mm	Total rain loss	63.4 mm
Actual water use by crop	380.3 mm	Moist deficit at harvest	60.8 mm
Potential water use by crop	380.3 mm	Actual irrigation requirement	60.8 mm
Efficiency irrigation schedule	- %	Efficiency rain	83.4 %
Deficiency irrigation schedule	0.0 %		

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------





2. Irrigation schedule option - output

Rainfed conditions

$$ET_{green} (irr = 0) = ET_{tot} (irr = 0)$$

$$ET_{blue} (irr = 0) = 0$$

Irrigated conditions

$$ET_{green} (irr = 1) = ET_{green} (irr = 0)$$

$$ET_{blue} (irr = 1) = ET_{tot} (irr = 1) - ET_{green} (irr = 0)$$

Rainfed scenario (*irr = 0*)

CROPWAT options

Non-rice crop scheduling |

Scheduling criteria for non-rice crops

Irrigation timing

No irrigation (rainfed)

No irrigation (only rainfall)

Irrigation application

Refill soil to field capacity

Irrigation efficiency

Irrigation efficiency: 70 %

Save as default Reset to FAO defaults OK Cancel Help

Irrigated scenario (*irr = 1*)

CROPWAT options

Non-rice crop scheduling |

Scheduling criteria for non-rice crops

Irrigation timing

Irrigate at critical depletion

Irrigation at 100 % critical depletion

Irrigation application

Refill soil to field capacity

Irrigation efficiency

Irrigation efficiency: 70 %

Save as default Reset to FAO defaults OK Cancel Help





2. Irrigation schedule option – output – Medium soil

Rainfed scenario ($irr = 0$)

Crop irrigation schedule										
ETo station	GRONINGEN-AP-EELD	Crop	Sugarbeet	Planting date	10/04	Yield red.				
Rain station	GRONINGEN-AP-EELD	Soil	Medium (loam)	Harvest date	06/10	0.0 %				
Table format				Timing: No irrigation (rainfed)						
<input type="radio"/> Irrigation schedule <input checked="" type="radio"/> Daily soil moisture balance				Application: - Field eff. 70 %						
Date	Day	Stage	Rain	Ks	Eta	Depl	Net Irr	Deficit	Loss	Gr. Irr
			mm	fract.	mm/day	%	mm	mm	mm	mm
29 Sep	173	End	0.0	1.00	1.3	23	0.0	66.3	0.0	0.0
30 Sep	174	End	0.0	1.00	1.3	23	0.0	67.6	0.0	0.0
1 Oct	175	End	0.0	1.00	0.9	24	0.0	68.5	0.0	0.0
2 Oct	176	End	0.0	1.00	0.9	24	0.0	69.4	0.0	0.0
3 Oct	177	End	11.4	1.00	0.9	20	0.0	58.9	0.0	0.0
4 Oct	178	End	0.0	1.00	0.9	21	0.0	59.9	0.0	0.0
5 Oct	179	End	0.0	1.00	0.9	21	0.0	60.8	0.0	0.0
6 Oct	End	End	0.0	1.00	0.0	21				

Totals

Total gross irrigation	0.0	mm	Total rainfall	383.0	mm
Total net irrigation	0.0	mm	Effective rainfall	319.5	mm
Total irrigation losses	0.0	mm	Total rain loss	63.4	mm
Actual water use by crop	380.3	mm	Moist deficit at harvest	60.8	mm
Potential water use by crop	380.3	mm	Actual irrigation requirement	60.8	mm
Efficiency irrigation schedule	-	%	Efficiency rain	83.4	%
Deficiency irrigation schedule	0.0	%			

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------

Rainfed conditions

$$ET_{green} (irr = 0) = ET_{tot} (irr = 0)$$

$$ET_{blue} (irr = 0) = 0$$





2. Irrigation schedule option – output – Medium soil

Irrigated scenario ($irr = 1$)

Crop irrigation schedule		Crop		Planting date		Yield red.				
ET ₀ station	GRONINGEN-AP-EELD	Crop	Sugarbeet	10/04		Harvest date	06/10			
Rain station	GRONINGEN-AP-EELD	Soil	Medium (loam)				0.0 %			
Table format		Timing: Irrigate at critical depletion								
<input type="radio"/> Irrigation schedule <input checked="" type="radio"/> Daily soil moisture balance		Application: Refill soil to field capacity Field eff. 70 %								
Date	Day	Stage	Rain	K _s	Eta	Depl.	Net Irr	Deficit	Loss	Gr. Irr
			mm	fract.	mm/day	%	mm	mm	mm	mm
29 Sep	173	End	0.0	1.00	1.3	23	0.0	66.3	0.0	0.0
30 Sep	174	End	0.0	1.00	1.3	23	0.0	67.6	0.0	0.0
1 Oct	175	End	0.0	1.00	0.9	24	0.0	68.5	0.0	0.0
2 Oct	176	End	0.0	1.00	0.9	24	0.0	69.4	0.0	0.0
3 Oct	177	End	11.4	1.00	0.9	20	0.0	58.9	0.0	0.0
4 Oct	178	End	0.0	1.00	0.9	21	0.0	59.9	0.0	0.0
5 Oct	179	End	0.0	1.00	0.9	21	0.0	60.8	0.0	0.0
6 Oct	End	End	0.0	1.00	0.0	21				

Totals

Total gross irrigation	0.0 mm	Total rainfall	383.0 mm
Total net irrigation	0.0 mm	Effective rainfall	319.5 mm
Total irrigation losses	0.0 mm	Total rain loss	63.4 mm
Actual water use by crop	380.3 mm	Moist deficit at harvest	60.8 mm
Potential water use by crop	380.3 mm	Actual irrigation requirement	60.8 mm
Efficiency irrigation schedule	- %	Efficiency rain	83.4 %
Deficiency irrigation schedule	0.0 %		

Yield reductions

Stagelabel	A	B	C	D	Season
------------	---	---	---	---	--------

Irrigated conditions

$$ET_{green} (irr = 1) = ET_{green} (irr = 0)$$

$$ET_{blue} (irr = 1) = ET_{tot} (irr = 1) - ET_{green} (irr = 0)$$





2. Irrigation schedule option - output – Soil comparison

Totals		Total gross irrigation	110.4	mm	Total rainfall	383.0	mm
		Total net irrigation	77.2	mm	Effective rainfall	300.0	mm
		Total irrigation losses	0.0	mm	Total rain loss	83.0	mm
		Actual water use by crop	380.3	mm	Moist deficit at harvest	3.1	mm
		Potential water use by crop	380.3	mm	Actual irrigation requirement	80.4	mm
		Efficiency irrigation schedule	100.0	%	Efficiency rain	78.3	%
		Deficiency irrigation schedule	0.0	%			
Totals		Total gross irrigation	0.0	mm	Total rainfall	383.0	mm
		Total net irrigation	0.0	mm	Effective rainfall	319.5	mm
		Total irrigation losses	0.0	mm	Total rain loss	63.4	mm
		Actual water use by crop	380.3	mm	Moist deficit at harvest	60.8	mm
		Potential water use by crop	380.3	mm	Actual irrigation requirement	60.8	mm
		Efficiency irrigation schedule	-	%	Efficiency rain	83.4	%
		Deficiency irrigation schedule	0.0	%			
Totals		Total gross irrigation	0.0	mm	Total rainfall	383.0	mm
		Total net irrigation	0.0	mm	Effective rainfall	319.5	mm
		Total irrigation losses	0.0	mm	Total rain loss	63.4	mm
		Actual water use by crop	380.3	mm	Moist deficit at harvest	60.8	mm
		Potential water use by crop	380.3	mm	Actual irrigation requirement	60.8	mm
		Efficiency irrigation schedule	-	%	Efficiency rain	83.4	%
		Deficiency irrigation schedule	0.0	%			





2. Irrigation schedule option - output – Soil comparison

CROPWAT option	ET_{green}	ET_{blue}	ET	CWU_{green}	CWU_{blue}	CWU	Y^*	$WF_{proc,green}$	$WF_{proc,blue}$	WF_{proc}
	mm/period			m^3/ha			ton/ha	m^3/ton		
Light	336	44	380	3364	440	3803	59	57	7	64
Medium	380	0	380	3803	0	3803	59	64	0	64
Heavy	380	0	380	3803	0	3803	59	64	0	64

*IRS (1999-2006) for Northern Netherlands





Water footprint estimation

1. CWR option

ET_g	ET_b	ET	CWU_g	CWU_b	CWU	Y^*	$WF_{proc,green}$	$WF_{proc,blue}$	WF_{proc}
mm/period			m³/ha			ton/ha	m³/ton		
282	100	381	2816	995	3811	59	48	17	65

*IRS (1999-2006) for Northern Netherlands

2. Irrigation schedule option

CROPWAT option	ET_{green}	ET_{blue}	ET	CWU_{green}	CWU_{blue}	CWU	Y^*	$WF_{proc,green}$	$WF_{proc,blue}$	WF_{proc}
	mm/period			m³/ha			ton/ha	m³/ton		
Light	336	44	380	3364	440	3803	59	57	7	64
Medium	380	0	380	3803	0	3803	59	64	0	64
Heavy	380	0	380	3803	0	3803	59	64	0	64

*IRS (1999-2006) for Northern Netherlands

Average irrigated area varies from 1 to 19% (IIRB, 2004)



Grey water footprint

- volume of polluted freshwater that associates with the production of a product in its full supply-chain.
- calculated as the volume of water that is required to assimilate pollutants based on ambient water quality standards.

Grey water footprint of a crop



$$\blacktriangleright \text{WF}_{\text{proc,grey}} \text{ (m}^3\text{)} = \frac{L}{(c_{\max} - c_{\text{nat}})}$$

$$\blacktriangleright \text{WF}_{\text{proc,grey}} \text{ (m}^3/\text{ton)} = \frac{L / (c_{\max} - c_{\text{nat}})}{\text{Prod}}$$

L – Load of pollutants entering the water system (g/day)

c_{\max} – Ambient water quality standard for the pollutant considered (g/m³)

c_{nat} – Natural concentration in the receiving water body (g/m³)

Prod – Production (ton/yr)



Grey water footprint

Source: Fertistat (FAO, 2009c)

 FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS
helping to build a world without hunger

Plant Production and Protection Division

FertiStat Fertilizer Use Statistics



[English](#) | [Français](#) | [Español](#)

<ul style="list-style-type: none">■ Home■ Fertilizer Use Statistics■ Download dataset■ Publications■ Related Sites ■ Contact	<p>Fertilizer Use Statistics</p> <p>This page provides 3 criteria for the selection of fertilizer use statistics. You may select more than one country by holding down the Ctrl key and clicking on the countries. FertiStat is still growing - if there is no data for the country-crop-year combination selected, please retry by dropping one of the selection criteria.</p> <p>Country</p> <p>Moldova, Republic of Morocco Myanmar Netherlands</p> <p>Commodity</p> <p>Beet</p> <p>Year</p> <p>All years</p> <p><input type="button" value="Search!"/> <input type="button" value="Reset"/></p>
--	--

[Contact FertiStat](#)

© FAO 2007



Grey water footprint

Source: Fertistat (FAO, 2009c)

Grey component of the process water footprint for sugarbeet within the Northern Netherlands

Average fertilizer application rate*	Area**	Total fertilizer applied	Nitrogen leached to the water bodies 10%	US EPA (2009)	Total WF _{proc,gray} ; sugar beet	Production**	WF _{proc,gray} ; sugar beet
kg/ha	ha	ton/year	ton/year	mg/l	10 ⁶ m ³ /year	ton	m ³ /ton
108	10887	1176	118	10	12	647176	18

*Fertistat (FAO, 2009c)

**IRS (1999-2006) for Northern Netherlands

References

- Allen, R.G., Pereira, L.S., Raes, D. and Smith, M. (1998)** Crop evapotranspiration - Guidelines for computing crop water requirements - FAO Irrigation and drainage paper 56. Food and Agriculture Organization. Rome. Available from: <http://www.fao.org/docrep/X0490E/X0490E00.htm>
- Chapagain, A.K. and Hoekstra, A.Y. (2004)** Water footprints of nations. Value of Water Research Report Series No. 16, UNESCO-IHE, Delft, the Netherlands. Available from: <http://www.waterfootprint.org/?page=files/Research%20data>
- EPA (2009)** Drinking Water Contaminants. US Environmental Protection Agency. Available from: <http://www.epa.gov/safewater/contaminants/index.html>
- FAO (2003)** Technical Conversion Factors for Agricultural Commodities. Food and Agriculture Organization of the United Nations, Rome. Online available: www.fao.org/WAICENT/FAOINFO/ECONOMIC/ESS/pdf/tcf.pdf
- FAO (2005)** New LocClim 1.10 database. Food and Agriculture Organization of the United Nations, Rome, Available from: http://www.fao.org/nr/climpag/pub/en3_051002_en.asp
- FAO (2006)** CLIMWAT Database. Food and Agriculture Organization, Rome. Available from: http://www.fao.org/nr/water/infores_databases_climwat.html
- FAO (2009a)** CROPWAT Model. Food and Agriculture Organization, Rome. Available from: http://www.fao.org/nr/water/infores_databases_cropwat.html
- FAO (2009b)** FAOSTAT Database. Food and Agriculture Organization. Rome. Available from: <http://faostat.fao.org>
- FAO (2009c)** FERTISTAT Database. Food and Agriculture Organization, Rome. Available from: http://www.fao.org/ag/agl/fertistat/fst_fubc_en.asp
- Hoekstra, A.Y. and Chapagain, A.K. (2008)** Globalization of water: Sharing the planet's freshwater resources. Blackwell Publishing. Oxford, UK.
- Monfreda, C., Ramankutty, N. and Foley, J.A. (2008)** Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, Global Biogeochemical Cycles, Vol. 22, GB1022, doi:10.1029/2007GB002947. Available from: <http://www.geog.mcgill.ca/~nramankutty/Datasets/Datasets.html>